PROBABILITY
AND
MATHEMATICAL STATISTICS
Vol. 0, Fasc. 0 (0000), pp. 000–000

arXiv:1704.01563v1 [math.PR] 5 Apr 2017

ON EXTREMAL INDEX OF MAX-STABLE STATIONARY PROCESSES

BY

KRZYSZTOF DĘBICKI (WROCŁAW) AND ENKELEJD HASHORVA (LAUSANNE)

To Tomasz Rolski,
thankful for all the support and ideas you shared with us!

Abstract. In this contribution we discuss the relation between Pickands-type constants defined for certain Brown-Resnick stationary process $W(t), t \in \mathbb{R}$ as

$$\mathcal{H}_W^\delta = \lim_{T\to\infty} T^{-1}\mathbb{E}\left\{\sup_{t\in\delta\mathbb{Z}\cap[0,T]} e^{W(t)}\right\},\ \delta \geqslant 0$$

(set $0\mathbb{Z} = \mathbb{R}$ if $\delta = 0$) and the extremal index of the associated max-stable stationary process ξ_W. We derive several new formulas and obtain lower bounds for $\mathcal{H}_W^\delta$ if W is a Gaussian or a Lévy process. As a by-product we show an interesting relation between Pickands constants and lower tail probabilities for fractional Brownian motions.

2000 AMS Mathematics Subject Classification: Primary: 60G15; Secondary: 60G70.

Key words and phrases: Extremal index, mean cluster index, Pickands constant, M3 representation, Brown-Resnick stationary, max-stable process, Gaussian process, Lévy process.

1. INTRODUCTION

The motivation for this contribution comes from the importance and the intriguing properties of the classical Pickands constants $\mathcal{H}_W^\delta$, which are defined for

any $\delta \geqslant 0$ by (interpret $0\mathbb{Z}$ as $\mathbb{R}$)

$$\mathcal{H}_W^\delta = \lim_{T\to\infty} \frac{1}{T}\mathbb{E}\left\{\sup_{t\in\delta\mathbb{Z}\cap[0,T]} e^{W(t)}\right\}, \tag{1.1}$$

where

$$W(t) = \sqrt{2}B_\alpha(t) - |t|^\alpha, \quad t \in \mathbb{R},$$

with B_α a standard fractional Brownian motion with Hurst index $\alpha \in (0, 2]$, that is a centered Gaussian process with stationary increments and variance function $\mathrm{Var}(B_\alpha(t)) = |t|^\alpha, t \in \mathbb{R}$.

It is well-known (but not trivial to prove) that $\mathcal{H}_W^\delta$ is finite and positive for any $\delta \geqslant 0$. The only values known for $\mathcal{H}_W^\delta$ are for $\delta = 0$ and $\alpha \in \{1, 2\}$, see e.g., [41, 42]. Suprisingly, Pickands and related constant appear in numerous unrelated asymptotic problems, see e.g., the recent papers [17, 25, 26, 15].

The contribution [19] derived a new formula for Pickands constants, which in fact indicates a direct connection between those contants and max-stable stationary processes, see [11]. The definition of $\mathcal{H}_W^\delta$ in (1.1) is extended in [11] for some general process W, provided that it defines a max-stable and stationary process. More precisely, assume throughout in the sequel that

$$W(t) = B(t) - \ln \mathbb{E}\left\{e^{B(t)}\right\}, \quad t \in \mathbb{R}, \tag{1.2}$$

where $B(t), t \in \mathbb{R}$ is a random process on the space D of càdlàg functions $f : \mathbb{R} \to \mathbb{R}$ with

$$B(0) = 0, \quad \mathbb{E}\left\{e^{B(t)}\right\} < \infty, \quad t \in \mathbb{R}. \tag{1.3}$$

Hence $X(t) = e^{W(t)}$ satisfies $X(0) = 1$ almost surely, and $\mathbb{E}\{X(t)\} = 1, t \in \mathbb{R}$. If $\Pi = \sum_{i=1}^\infty \varepsilon_{P_i}$ is a Poisson point process (PPP) with intensity $x^{-2}dx$ on $(0, \infty)$, and $X_i = e^{W_i}, i \geqslant 1$ are independent copies of the random process $X = e^W$ being independent of Π, then the random process ξ_W defined by

$$\xi_W(t) = \max_{i\geqslant 1} P_i X_i(t) = \max_{i\geqslant 1} P_i e^{W_i(t)}, \quad t \in \mathbb{R} \tag{1.4}$$

has unit Fréchet marginals and is max-stable. Here ε_x denotes the unit Dirac measure at $x \in \mathbb{R}$.

Adopting the definition in [30], we shall refer to W as the *Brown-Resnick stationary process* whenever the associated max-stable process ξ_W is stationary. Note that stationarity of ξ_W means that $\{\xi_W(t), t \in \mathbb{R}\}$ and $\{\xi_W(t+h), t \in \mathbb{R}\}$ have the same distribution for any $h \in \mathbb{R}$.

In the sequel, for the case $\delta = 0$ we shall assume that

$$\mathbb{E}\left\{\sup_{t\in K} e^{W(t)}\right\} < \infty$$

for any compact $K \subset \mathbb{R}$. A direct consequence of stationarity of ξ_W and the fact that for any $t_1, \dots, t_n \in \mathbb{R}$ and $x_1, \dots, x_n > 0$, see e.g., [18, 39]

$$\mathbb{P}\left\{\xi_W(t_i) \leqslant x_i, \forall i \leqslant n\right\} = e^{-\mathbb{E}\left\{\max_{1\leqslant i\leqslant n}\left(e^{W(t_i)}/x_i\right)\right\}} \tag{1.5}$$

is that, for any $b \geqslant 0, \delta \geqslant 0, T > 0$ we have

$$H^\delta_W([0,T]) \;:=\; \mathbb{E}\left\{\sup_{t\in\delta\mathbb{Z}\cap[0,T]} e^{W(t)}\right\} = \mathbb{E}\left\{\sup_{t\in\delta\mathbb{Z}\cap[b,b+T]} e^{W(t)}\right\}.$$

Consequently, $\mathcal{H}^\delta_W$ defined in (1.1) exists and is given by (see [11])

$$\mathcal{H}^\delta_W = \inf_{T>0} \frac{1}{T} H^\delta_W([0,T]) \in [0,\infty). \tag{1.6}$$

Note that if $\delta > 0$, then (1.6) implies that

$$\mathcal{H}^\delta_W \leqslant \frac{H^\delta_W([0,\delta-\varepsilon])}{\delta-\varepsilon} = \frac{1}{\delta-\varepsilon}$$

for any $\varepsilon \in (0,\delta)$, hence letting ε tend to 0 yields $\mathcal{H}^\delta_W \in [0,\delta]$.

Interestingly, $\mathcal{H}^\delta_W$ is related to the extremal index of the stationary process

$$\xi^\delta_W(t) = \xi_W(\delta t), \quad t \in \mathbb{Z}, \delta > 0,$$

where we set $\xi^\delta_W(t) = \xi_W(t)$ if $\delta = 0$. Indeed, by (1.5)

$$\begin{aligned}\lim_{T\to\infty} \mathbb{P}\left\{\max_{i\in\delta\mathbb{Z}\cap[0,T]} \xi_W(t) \leqslant Tx\right\} &= e^{-\lim_{T\to\infty}\mathbb{E}\left\{\max_{i\in\delta\mathbb{Z}\cap[0,T]}\left(e^{W(i)}/T\right)\right\}\frac{1}{x}} \\ &= \left(e^{-\frac{1}{x}}\right)^{\mathcal{H}^\delta_W}, \quad x > 0. \end{aligned} \tag{1.7}$$

Thus the Fréchet limit result in (1.7), which is already shown in [50] (see also [10][Proposition 3.1] and [18]) states that the extremal index of the stationary process $\xi_W^\delta(t), t \in \mathbb{Z}$ is given for any $\delta > 0$ by

$$\theta_W^\delta = \delta \mathcal{H}_W^\delta \in [0,1]. \tag{1.8}$$

Clearly, the constant $\mathcal{H}_W^\delta$ is positive if and only if the extremal index θ_W^δ of the stationary process ξ_W^δ is positive.

Numerous papers in the literature have discussed the calculation and estimation of extremal index of stationary processes, see e.g., the recent articles [46, 10, 38, 35, 33, 21] and the references therein.

The primary goal of this contribution is to study Pickands type constants $\mathcal{H}_W^\delta$ by exploring the properties of the extremal index θ_W^δ. In particular, we are interested in establishing tractable conditions that guarantee the positivity of $\mathcal{H}_W^\delta$.

By our assumptions it is clear that ξ_W^δ is stationary and *jointly regularly varying*, hence in view of [5, Theorem 2.1] (see also [29]), there exists the so-called *tail process*

$$Y^\delta(i), \quad i \in \mathbb{Z}$$

of the stationary process X, which was introduced in [5]. It turns out that for any $m \leqslant n, m, n \in \mathbb{Z}$ we have the following stochastic representation

$$(Y^\delta(m), \ldots, Y^\delta(n)) \stackrel{d}{=} (\mathcal{P}X^\delta(m), \ldots, \mathcal{P}X^\delta(n)), \tag{1.9}$$

$X^\delta(i) := e^{W(\delta i)}, i \in \mathbb{Z}$, with $\mathcal{P}$ a unit Pareto random variable with survival function $1/x, x > 1$ being independent of the process X.

Under the *finite mean cluster size condition* (see below Condition 2.1) and condition $\mathcal{A}(a_n)$, see [5, 4, 32], it follows that θ_W^δ is positive, see the seminal contribution [5].

We shall show the positivity of the extremal index under a weaker condition, namely supposing that

$$\lim_{|z| \to \infty, z \in \mathbb{Z}} W(z\delta) = -\infty \tag{1.10}$$

holds almost surely for $\delta \in (0, \infty)$. In our derivations the next simple result is crucial:

LEMMA 1.1. *If $r_n, n \geqslant 1$ are positive integers such that*

$$\lim_{n\to\infty} r_n = \lim_{n\to\infty} n/r_n = \infty,$$

then for any $\delta \in (0,\infty)$ we have

$$\widetilde{\theta^{\delta}_W} := \lim_{n\to\infty} \frac{n}{r_n}\mathbb{P}\left\{\max_{i\in\{0,\delta,\ldots,\delta r_n\}} \xi_W(t) > n\right\} = \theta^{\delta}_W \in [0,\infty). \tag{1.11}$$

In the next section we shall show that the new expression for the extremal index in (1.11) is positive under (1.10). Exploiting the explicit form of the *tail process* we shall derive several new interesting formulas for $\mathcal{H}^{\delta}_W$.

Brief outline of the rest of the paper: Section 2 displays our main results which establish the positivity of the Pickands-type constants and some new formulas. In Section 3 we shall discuss the connection with mixed moving maxima (M3) representation of Brown-Resnick processes. Then we derive some explicit lower bounds for $\mathcal{H}^{\delta}_W$ in case that B in (1.2) is a Gaussian or a Lévy process and then discuss the relation between $\mathcal{H}^{0}_W$ and the *mean cluster index*. Further, we shall show that the classical Pickands constants are related to a small ball problem. All the proofs are relegated to Section 4.

2. MAIN RESULTS

We keep the same setup as in the Introduction and denote additionally by $\mathcal{E}$ a unit exponential random variable which is independent of everything else. According to [5] a candidate for the extremal index is given by the following formula

$$\widehat{\theta^{\delta}_W} = \lim_{m\to\infty} \mathbb{P}\left\{\max_{1\leqslant i\leqslant m} Y^{\delta}(i) \leqslant 1\right\}, \tag{2.1}$$

where $Y^{\delta}(i), i \in \mathbb{Z}$ is the *tail process* of ξ^{δ}_W, see [5]. As in the aforementioned paper we shall impose the *finite mean cluster size condition* of [5, Condition 4.1]:

CONDITION 2.1. *Given $\delta > 0$, there exists a sequence of positive integers $r_n, n \in \mathbb{N}$ satisfying $\lim_{n\to\infty} r_n/n = 1/\lim_{n\to\infty} r_n = 0$ such that*

$$\lim_{m\to\infty} \limsup_{n\to\infty} \mathbb{P}\left\{\max_{m\leqslant|k|\leqslant r_n} \xi_W(k\delta) > nx \middle| \xi_W(0) > nx\right\} = 0 \tag{2.2}$$

holds for any $x > 0$.

In view of [5, Proposition 4.2] we have that $\widehat{\theta^{\delta}_W} > 0$ follows from Condition 2.1. Our main result below establishes new formulas for $\mathcal{H}^{\delta}_W$.
Moreover, from the above mentioned reference, Condition 2.1 together with well-known $\mathcal{A}(a_n)$ conditions of Hsing and Davis implies that the candidate of extremal index is equal to the extremal index, i.e., $\widehat{\theta^{\delta}_W} = \theta^{\delta}_W > 0$. It is well-known that $\mathcal{A}(a_n)$ is implied by the strong mixing of ξ^{δ}_W. However, our results derived below do not require strong mixing, but just mixing of ξ^{δ}_W.

THEOREM 2.1. *Let* $X(t) = e^{W(t)}, t \in \mathbb{R}$ *with* W *as in* (1.2) *be such that* (1.3) *holds and* $\xi_W(t), t \in \mathbb{R}$ *is max-stable and stationary. We have that* (1.10) *holds for* $\delta > 0$ *if and only if Condition* (2.1) *holds. Moreover, if* (1.10) *holds for* $\delta > 0$, *then*

$$\mathcal{H}^{\delta}_W = \frac{1}{\delta}\mathbb{P}\left\{\sup_{i<0} W^{\delta}(i) < 0 = \sup_{i\in\mathbb{Z}} W^{\delta}(i)\right\} \tag{2.3}$$

$$= \frac{1}{\delta}\mathbb{P}\left\{\sup_{i\geqslant 1}\big(\mathcal{E} + W^{\delta}(i)\big) \leqslant 0\right\} \tag{2.4}$$

$$= \frac{1}{\delta}\Big[\mathbb{E}\left\{\sup_{i\geqslant 0} e^{W^{\delta}(i)}\right\} - \mathbb{E}\left\{\sup_{i\geqslant 1} e^{W^{\delta}(i)}\right\}\Big] \in (0, 1/\delta), \tag{2.5}$$

where $W^{\delta}(t) = W(t\delta), t \in \mathbb{Z}$ *and* $\mathcal{E}$ *is a unit exponential random variable independent of* W.

REMARK 2.1. *a) If* $\mathbb{P}\left\{W^{\delta}(i) = 0\right\} = 0$ *for any negative integer* i, *then*

$$\mathbb{P}\left\{\sup_{-m\leqslant i<0} W^{\delta}(i) < 0 = \sup_{-m\leqslant j\leqslant m} W^{\delta}(j)\right\} = \mathbb{P}\left\{\sup_{-m\leqslant j\leqslant m} W^{\delta}(j) = 0\right\}$$

holds for any integer $m > 1$. *Consequently, by* (2.3) *we have*

$$\mathcal{H}^{\delta}_W = \frac{1}{\delta}\lim_{m\to\infty}\mathbb{P}\left\{\sup_{-m\leqslant i<0} W^{\delta}(i) < 0 = \sup_{-m\leqslant j\leqslant m} W^{\delta}(j)\right\}$$

$$= \frac{1}{\delta}\mathbb{P}\left\{\sup_{i\in\mathbb{Z}} W^{\delta}(i) = 0\right\} > 0, \tag{2.6}$$

which has been shown in [19] for the case B *is a standard fractional Brownian motion. The assumption* $W(0) = 0$ *can be removed, see [27].*

b) Above we assumed that ξ_W has càdlàg sample paths in order to define H^0_W. For the results of Theorem 2.1, this assumption is not needed.

c) In [11] it is shown that under the assumptions of Theorem 2.1 we have

$$\mathcal{H}^\delta_W = \mathbb{E}\left\{\frac{\sup_{t\in\delta\mathbb{Z}} e^{W(t)}}{\delta \sum\limits_{t\in\delta\mathbb{Z}} e^{W(t)}}\right\}. \tag{2.7}$$

According to (2.5), for calculation of $\mathcal{H}^\delta_W$ it suffices to know $W(t), t\in\delta\mathbb{Z}, t>0$, i.e., only the values of W for positive t matter. This is not the case for the formula (2.7). Both (2.7) and (2.5) are given in terms of expectations and not as limits, which is a great advantage for simulations. To this end, we mention that simulation of Pickands constants has been the topic of many contributions, see e.g., [9, 36, 19].

d) If $X(t)=e^{W(t)}, t\in\mathbb{R}$ is Brown-Resnick stationary, i.e., the associated max-stable process with ζ_W is max-stable and stationary, then the time reversed process $V(t)=W(-t), t\in\mathbb{R}$ also defines a Brown-Resnick stationary processes. Moreover, for any $\delta\geqslant 0$

$$\mathcal{H}^\delta_W = \mathcal{H}^\delta_V.$$

Consequently the formulas in Theorem 2.1 can be stated with V instead of W, for instance we have

$$\begin{aligned}\mathcal{H}^\delta_W &= \frac{1}{\delta}\mathbb{P}\left\{\sup_{i\leqslant -1}\left(\mathcal{E}+W^\delta(i)\right)\leqslant 0\right\}\\ &= \frac{1}{\delta}\mathbb{P}\left\{W^\delta(i)<0, i\in\mathbb{N}, W^\delta(i)\leqslant 0, i\in\mathbb{Z}\right\}.\end{aligned} \tag{2.8}$$

e) If $W(t)=\sqrt{2}tL-t^2$ with L an $N(0,1)$ random variable with distribution Φ and probability density function φ, by (2.4) we have

$$\begin{aligned}\mathcal{H}^\delta_W &= \frac{1}{\delta}\int_0^\infty \mathbb{P}\left\{\mathcal{E}+\sup_{i\geqslant 1}\left(\sqrt{2}\delta i b-(\delta i)^2\right)\leqslant 0\right\}\varphi(b)db\\ &= \frac{1}{\delta}\int_{-\delta/\sqrt{2}}^{\delta/\sqrt{2}}\varphi(b)db = \frac{1}{\delta}\left[\Phi(\delta/\sqrt{2})-\Phi(-\delta/\sqrt{2})\right]\end{aligned} \tag{2.9}$$

holds for any $\delta > 0$. *Consequently, letting* $\delta \to 0$ *we obtain the well-known result*

$$\mathcal{H}_W^0 = \sqrt{2}\varphi(0) = \frac{1}{\sqrt{\pi}}.$$

A canonical example for W with representation (1.2) is the case when B is a centered Gaussian process with stationary increments, continuous sample paths, and variance function σ^2. Then the max-stable process ξ_W is stationary, see [40]. Using a direct argument, we establish in the next theorem the positivity of $\mathcal{H}_W^0$.

THEOREM 2.2. *If*

$$\liminf_{t\to\infty} \frac{\sigma^2(t)}{\ln t} > 8, \tag{2.10}$$

then $\mathcal{H}_W^0 > 0$.

Since (2.10) implies (1.10), see Corollary 2.4 in [34] or [30], then using $\mathcal{H}_W^0 \geqslant \mathcal{H}_W^\delta$ for any $\delta > 0$ we immediately establish the positivity of $\mathcal{H}_W^0$.

Indeed, the positivity of $\mathcal{H}_W^0$ is crucial for the study of extremes of Gaussian processes. Condition (2.10) can be easily checked, for instance if $W(t) = \sqrt{2}B_\alpha(t) - |t|^\alpha$. Consequently, the classical Pickands constants $\mathcal{H}_W^\delta$ are positive for any $\delta \geqslant 0$. This fact is highly non-trivial; after announced in Pickands' pioneering work [41], correct proofs were obtained later by Pickands himself, and in [7, 43], see for instance Theorem B3 in [8]. We note in passing that under general conditions on σ^2 the positivity of $\mathcal{H}_W^0$ is established in [13].

Apart from the alternative proof for the positiveness of the original Pickands constants, Theorem 2.1 extends to non-Gaussian processes W. For the above Gaussian setup, direct calculations show the positivity of $\mathcal{H}_W^\delta$ under a slightly weaker condition than (2.10).

3. DISCUSSIONS & EXTENSIONS

3.1. Relation with lower tail probabilities. For the classical case of Piterbarg constants $\mathcal{H}_{B_\alpha}$, i.e., for $W(t) = \sqrt{2}B_\alpha(t) - |t|^\alpha$, $t \in \mathbb{R}, \alpha \in (0, 2]$ we show below that (2.6) implies a nice relation with a small ball problem.

PROPOSITION 3.1. *For any* $\alpha \in (0, 2]$ *we have*

$$\lim_{\eta \to 0} \eta^{-2/\alpha} \mathbb{P}\left\{\forall_{k \in \mathbb{Z} \setminus \{0\}} B_\alpha(1/k) \leqslant \eta\right\} = 2^{1/\alpha} \mathcal{H}_{B_\alpha}.$$

The above result strongly relates to the self-similarity property of fractional Brownian motion. In case of a general Gaussian W, we still have that ξ_W is stationary if W has stationary increments. However, fBm is the only centered Gaussian process with stationary increments being further self-similar. Hence, no obvious extensions of the above relation with lower tails can be derived for general W.

3.2. Non-Gaussian W. The classical Pickands constants are defined for $W(t) = \sqrt{2} B_\alpha(t) - |t|^\alpha$ with B_α a standard fBm with Hurst index $\alpha/2 \in (0, 1]$. The more general case where B_α is substituted by a centered Gaussian process with stationary increments is discussed in details in [13].

Our setup clearly allows for any random process W, not necessarily Gaussian, which is Brown-Resnick stationary. Along with the Gaussian case of W, the Lévy one has also been dealt already in the literature. In view of [23, 49], if $B(t), t \geqslant 0$ is a Lévy process such that

$$\Phi(1) < \infty, \quad \Phi(\theta) := \ln \mathbb{E}\left\{e^{\theta B(1)}\right\},$$

then $W(t) = B(t) - \Phi(1)t, t \geqslant 0$ is Brown-Resnick stationary, i.e., $\xi_W(t), t \geqslant 0$ is max-stable stationary with unit Gumbel marginals.

In [31] an important constant appears in the asymptotic analysis of the maximum of standardised increments of random walks, which in fact is the Pickands constant $\mathcal{H}^\delta_W, \delta > 0$ introduced here for W as above. In [31][Lemma 5.16] a new formula for $\mathcal{H}^\delta_W$ is derived, which is identical with our formula in (2.8). Another instance of the Pickands constant given by formula (2.3) is displayed in [44][Theorem 5.3]. With the notation of that theorem, we have for $\delta = 1$ that

$$W(i) = \sum_{j=1}^{i} A_i,$$

where A_i's are iid with the same distribution as $Z\mathbb{I}(U \leqslant e^{-\eta Z})$ for some $\eta > 0$ with U uniformly distributed on $(0, 1)$ being independent of Z which has some pdf

symmetric around 0.

Pickands constants appear also in the context of semi-min-stable processes, see [51]. In view of the aforementioned paper, several results derived here for max-stable processes are extendable to semi-min-stable processes.

3.3. Finite Mean Cluster Size Condition. As noted in [45], Condition 2.1 is implied by the so-called *short-lasting exceedance condition* given below:

CONDITION 3.1. *Given $\delta > 0$, there exists a sequence of integers $r_n, n \in \mathbb{N}$ satisfying $\lim_{n\to\infty} r_n/n = 1/\lim_{n\to\infty} r_n = 0$ such that*

$$\lim_{m\to\infty} \limsup_{n\to\infty} \sum_{k=m}^{r_n} \mathbb{P}\left\{\xi_W(k\delta) > nx \middle| \xi_W(0) > nx\right\} = 0 \tag{3.1}$$

is valid for any $x > 0$.

This latter condition is a rephrasing of the so-called B condition, see e.g., [1, 12, 2], which was formulated by discretising the original Berman's condition, see [6]. Condition 3.1 is weaker than the $D'(xn)$ condition of Leadbetter as discussed in [22][Section 5.3.2].
Commonly, Condition (2.1) assumed for $x = 1$ is referred to as the *anti-clustering condition*, see e.g., [46, 47]. Clearly, the finite mean cluster size condition is stronger then the anti-clustering condition. The latter appears in various contexts related to extremes of stationary processes, see e.g., [3, 37, 46, 5, 47] and the references therein.

3.4. M3 Representation. Since we assume that ξ_W is max-stable stationary with càdlàg sample paths and W with representation (1.2) is such that B satisfies (1.3), then assuming the following almost sure convergence

$$W(t) \to -\infty \tag{3.2}$$

as $|t| \to \infty$ is equivalent with the fact that ξ_W possesses a *mixed moving maxima representation* (for short M3), see [20, Theorem 3] and [52]. More specifically,

under (3.2) we have the equality of finite dimensional distributions

$$\xi_W(t) \stackrel{d}{=} \max_{i\geqslant 1} P_i e^{F_i(t-T_i)}, \quad t \in \mathbb{R} \tag{3.3}$$

between rhs and lhs in (3.3), where the F_i's are independent copies of a measurable càdlàg process $F_W(t), t \in \mathbb{R}$ satisfying

$$\sup_{t\in\mathbb{R}} F_W(t) = F_W(0) = 0 \tag{3.4}$$

almost surely, and $\sum_{i=1}^{\infty} \varepsilon_{(P_i,T_i)}$ is a PPP in $(0,\infty)\times\mathbb{R}$ with intensity $C_W \cdot p^{-2}dp \cdot dt$ with

$$C_W = \left(\mathbb{E}\left\{\int_{\mathbb{R}} e^{F_W(t)}\, dt\right\}\right)^{-1} \in (0,\infty). \tag{3.5}$$

Moreover ξ_W^δ, the restriction of ξ_W on $\delta\mathbb{Z}$ possesses an M3 representation for any $\delta > 0$, see [11] for more details. Denote the corresponding constant in the intensity of this PPP by $C_W^\delta > 0$ (and thus C_W^0 is just C_W given in (3.5)).
In view of [11][Proposition 1], if $\xi_W^\delta, \delta > 0$ admits an M3 representation as mentioned above, then

$$\mathcal{H}_W^\delta = C_W^\delta, \tag{3.6}$$

provided that (1.10) holds. Hence Theorem 2.1 presents new formulas for C_W^δ. Note in passing that (3.6) has been shown in [40]. Therein it is proved that C_W^δ is given by the right-hand side of (2.6) assuming further that $W(t) = B(t) - \mathbb{E}\left\{e^{\ln B(t)}\right\}, t \in \mathbb{R}$ with B a centered Gaussian process with statioanry increments satisfying $W(0) = 0$ almost surely.
In view of [11][Theorem 1], if (1.10) holds, then we have

$$\mathcal{H}_W^\delta = \mathbb{E}\left\{\frac{M^\delta}{S^\delta}\right\} = C_W^\delta, \tag{3.7}$$

with $M^\delta := \max_{i\in\mathbb{Z}} e^{W(i\delta)}$ and $S^\delta := \delta\sum_{t\in\delta\mathbb{Z}} e^{W(t)}$. Thus $\mathcal{H}_W^\delta > 0$.
The representation of $\mathcal{H}_W^\delta$ as an expectation of the ratio M^δ/S^δ is crucial for its simulation. Such a representation has been initially shown in [19] for classical Pickands constants.

3.5. Lower Bounds. In Theorem 2.1 we present new formulas for $\mathcal{H}^{\delta}_{W}$, which in turn establish the positivity of $\mathcal{H}^{\delta}_{W}$ and thus the positivity for the extremal index of ξ^{δ}_{W}. If only the positivity of $\mathcal{H}^{\delta}_{W}$ is of primary interest, then the conditions of Theorem 2.1 can be relaxed. Next, we consider two important classes of processes for B that is centered Gaussian processes with stationary increments and Lévy processes. Results for the Lévy case has been already given in [11].

For particular values of δ, we show that it is possible to derive a positive lower bound for $\mathcal{H}^{\delta}_{W}$ and thus establishing the positivity of $\mathcal{H}^{\delta}_{W}$. Let $x_+ := \max(x, 0)$.

THEOREM 3.1. *i) Let $W(t) = B(t) - \sigma^2(t)/2$, $t \geqslant 0$, where $B(t)$ is a centered Gaussian processes with stationary increments and variance function σ^2 such that $\sigma(0) = 0$. Then for any $\delta > 0$*

$$\mathcal{H}^{\delta}_{W} \geqslant \frac{1}{\delta}\max\big(0, 1 - \sum_{k=1}^{\infty} e^{-\frac{\sigma^2(\delta k)}{8}}\big). \tag{3.8}$$

ii) Let $W(t) = B(t) - \Phi(1)t$, $t \geqslant 0$, where $B(t)$ is a Lévy process satisfying (3.1). *Then for any $\delta > 0$*

$$\mathcal{H}^{\delta}_{W} \geqslant \frac{1}{\delta}\frac{\max\big(0, 1 - 2e^{\left(\Phi(1/2) - \frac{1}{2}\Phi(1)\right)\delta}\big)}{1 - e^{\left(\Phi(1/2) - \frac{1}{2}\Phi(1)\right)\delta}}. \tag{3.9}$$

REMARK 3.1. *a) It follows from i) of Theorem 3.1 that if $\sigma(\delta k) \geqslant C(\delta k)^{\kappa/2}$ for all $k \in \mathbb{N}$ and some $\kappa > 0$, then*

$$\mathcal{H}^{\delta}_{W} \geqslant \frac{1}{\delta}\left(1 - \frac{1}{\delta}\frac{\Gamma(1/\kappa)}{\kappa\,(C^2/8)^{1/\kappa}}\right). \tag{3.10}$$

Since $\mathcal{H}^{0}_{W} \geqslant \mathcal{H}^{\delta}_{W}$ for any $\delta > 0$, then the above implies $\mathcal{H}^{0}_{W} > 0$.

b) If B is a Lévy process as in Theorem 3.1, ii), then (see the proof in Section 3)

$$\mathcal{H}^{0}_{W} \geqslant \frac{1}{8}[\Phi(1) - 2\Phi(1/2)] > 0. \tag{3.11}$$

3.6. Case $\delta = 0$. Since (1.7) holds also for $\delta = 0$ and $\mathcal{H}^{0}_{W} \geqslant \mathcal{H}^{\delta}_{W}$, then the extremal index of the continuous process ξ_W is

$$\widetilde{\theta}_W = \mathcal{H}^{0}_{W} \geqslant 0,$$

which is positive, provided that (1.10) holds. In the special case that $W(t) = \sqrt{2}B_\alpha(t) - |t|^\alpha$ we have that

$$\lim_{\delta \downarrow 0} \mathcal{H}_W^\delta = \mathcal{H}_W^0 =: \mathcal{H}_W, \tag{3.12}$$

hence for such W and for any $\alpha \in (0, 2]$

$$\widetilde{\theta}_W = \lim_{\delta \downarrow 0} \frac{\theta_W^\delta}{\delta}. \tag{3.13}$$

Recall that we denote by $\theta_W^\delta, \delta > 0$ the extremal index of ξ_W^δ. Using the terminology of [28] we refer to $\overline{H}_W$ defined by (assuming that the limit exists)

$$\lim_{\delta \downarrow 0} \frac{\theta_W^\delta}{\delta} = \lim_{\delta \downarrow 0} \mathcal{H}_W^\delta = \overline{H}_W$$

as the *mean cluster index of the process* W. Since for any $T > 0$ and $\delta > 0$

$$0 \leqslant \mathbb{E}\left\{ \sup_{t \in \delta\mathbb{Z} \cap [0,T]} e^{W(t)} \right\} =: \mathcal{H}_W^0([0, T]),$$

then clearly $\overline{H}_W \in [0, \mathcal{H}_W]$.
We show next that if ξ_W possesses an M3 representation, then $\overline{H}_W$ is positive.

PROPOSITION 3.2. *Suppose that ξ_W is max-stable and stationary with $W(0) = 0$. If ξ_W possesses an M3 representation and $\overline{H}_W$ exists, then*

$$\overline{H}_W \geqslant \mathbb{E}\left\{ \frac{\sup_{t \in \mathbb{R}} e^{W(t)}}{\eta \sum\limits_{t \in \eta\mathbb{Z}} e^{W(t)}} \right\} > 0 \tag{3.14}$$

holds for any $\eta > 0$.

REMARK 3.2. *a) In view of Theorems 2 and 3 in [11] we have for some general W as in (1.2), with B being Gaussian or Lévy process*

$$\mathcal{H}_W^0 = \mathbb{E}\left\{ \frac{\sup_{t \in \mathbb{R}} e^{W(t)}}{\eta \sum\limits_{t \in \eta\mathbb{Z}} e^{W(t)}} \right\} = \mathbb{E}\left\{ \frac{\sup_{t \in \mathbb{R}} e^{W(t)}}{\int\limits_{t \in \mathbb{R}} e^{W(t)}\, dt} \right\} \tag{3.15}$$

is valid for any $\eta > 0$. Consequently, under these conditions and the setup of Proposition 3.2

$$\mathcal{H}^0_W = \overline{H}_W. \tag{3.16}$$

b) If $W(t) = \sqrt{2}B_\alpha(t) - |t|^\alpha, t \in \mathbb{R}$, by (3.12) and (2.4) for any $\alpha \in (0, 2]$

$$\mathcal{H}^0_W \quad = \quad \overline{H}_W = \lim_{\delta \downarrow 0} \frac{1}{\delta} \mathbb{P}\left\{ \sup_{i \geqslant 1} (\mathcal{E} + W^\delta(i)) \leqslant 0 \right\}, \tag{3.17}$$

with $\mathcal{E}$ a unit exponential random variable independent of W. Expression (3.17) of the classical Pickands constant was initially derived in [1] for some general W, see also recent contribution [2]. In [28], Proposition 3 or the formula in [24][p.44] the classical Pickands constant is the limit of a cluster index.

4. PROOFS

Proof of Lemma 1.1: Since $\lim_{n\to\infty} r_n = \infty$, then by (1.7) and (1.8)

$$\lim_{n\to\infty} r_n^{-1} \mathbb{E}\left\{ \max_{i \in \{0,\delta,\ldots,\delta r_n\}} e^{W(i)} \right\} = \delta \mathcal{H}^\delta_W = \theta^\delta_W.$$

For any $n \in \mathbb{N}$ we have

$$\begin{aligned} & \frac{\mathbb{P}\left\{\max_{i\in\{0,\delta,\ldots,\delta r_n\}} \xi_W(i) > n\right\}}{r_n \mathbb{P}\left\{\xi_W(0) > n\right\}} \\ = \quad & \frac{\mathbb{P}\left\{\max_{i\in\{0,\delta,\ldots,\delta r_n\}} \xi_W(i) > n\right\}}{r_n[1 - e^{-1/n}]} \\ \sim \quad & n r_n^{-1}\Big[1 - \mathbb{P}\left\{ \max_{i \in \{0,\delta,\ldots,\delta r_n\}} \xi_W(i) \leqslant n \right\}\Big] \\ = \quad & n r_n^{-1}\big[1 - e^{-c_n/n}\big], \quad c_n := \mathbb{E}\left\{ \max_{i \in \{0,\delta,\ldots,\delta r_n\}} e^{W(i)} \right\}, \end{aligned}$$

where the last equality follows from (1.5). The assumption that $\lim_{n\to\infty} n/r_n = \infty$ and $\mathbb{E}\left\{e^{W(i)}\right\} = 1, i \in \delta\mathbb{Z}$ imply

$$\frac{c_n}{n} \leqslant \frac{1}{n} \mathbb{E}\left\{ \sum_{i \in \{0,\delta,\ldots,\delta r_n\}} e^{W(i)} \right\} = \frac{r_n + 1}{n} \to 0, \quad n \to \infty. \tag{4.1}$$

Consequently,

$$\frac{\mathbb{P}\left\{\max_{i\in\{0,\delta,\ldots,\delta r_n\}}\xi_W(i)>n\right\}}{r_n\mathbb{P}\left\{\xi_W(0)>n\right\}} \sim r_n^{-1}\mathbb{E}\left\{\max_{i\in\{0,\delta,\ldots,\delta r_n\}}e^{W(i)}\right\}\sim\theta_W^{\delta},\quad n\to\infty,$$

hence the claim follows. □

Proof of Theorem 2.1: We show first stochastic representation (1.9). Recall that $X(t)=e^{W(t)}$ and for $\delta>0$ we set

$$W^{\delta}(t)=W(\delta t),\quad X^{\delta}(t)=e^{W^{\delta}(t)},\quad t\in\mathbb{Z}.$$

By (1.5), the fact that $\mathbb{P}\{\xi_W(0)\leqslant x\}=e^{-1/x}, x>0$ and the assumption that $X(0)=1$ almost surely, for any $y_1,\ldots,y_n$ positive and $y_0>1$ we have

$$\begin{aligned}
&\mathbb{P}\left\{\xi^{\delta}_W(i)\leqslant Ty_i, i=0,\ldots,n\big|\xi^{\delta}_W(0)>T\right\}\\
&= \frac{1-\mathbb{P}\left\{\xi^{\delta}_W(0)\leqslant T,\xi^{\delta}_W(i)\leqslant Ty_i, i\in\{0,\ldots,n\}\right\}-[1-\mathbb{P}\left\{\xi^{\delta}_W(i)\leqslant Ty_i, i\in\{0,\ldots,n\}\right\}]}{\mathbb{P}\left\{\xi^{\delta}_W(0)>T\right\}}\\
&= \frac{1-e^{-\mathbb{E}\{\max\left(X^{\delta}(0),\max_{i\in\{1,\ldots,n\}}\frac{X^{\delta}(i)}{y_i}\right)\}\frac{1}{T}}-\left[1-e^{-\mathbb{E}\{\max_{i\in\{1,\ldots,n\}}\frac{X^{\delta}(i)}{y_i}\}\frac{1}{T}}\right]}{1-e^{-\frac{1}{T}}}\\
&\sim T\Big[1-\big[1-\tfrac{1}{T}\mathbb{E}\{\max\big(1,\max_{i\in\{0,\ldots,n\}}\frac{X^{\delta}(i)}{y_i}\big)\}\big]-\Big(1-\big[1-\tfrac{1}{T}\mathbb{E}\{\max_{i\in\{0,\ldots,n\}}\frac{X^{\delta}(i)}{y_i}\}\big]\Big)\Big]\\
&\to \mathbb{E}\left\{\left(1-\max_{i\in\{0,\ldots,n\}}\frac{X^{\delta}(i)}{y_i}\right)_+\right\},\quad T\to\infty\\
&= \mathbb{P}\left\{\mathcal{P}\leqslant y_0,\mathcal{P}X^{\delta}(i)\leqslant y_i,\forall i\in\{1,\ldots,n\}\right\},
\end{aligned}$$

where $\mathcal{P}$ is a unit Pareto random variable with survival function $1/s, s>1$ independent of the process X. Hence the claim in (1.9) follows by [5][Theorem 2.1 (ii)]. Next by the above derivations for any sequence of integers $r_n>m\in\mathbb{N}$ for any $x>0$ (recall $X^{\delta}(0)=1$ almost surely) we have

$$\begin{aligned}
&1-\mathbb{P}\left\{\max_{m\leqslant|i|\leqslant r_n}\xi^{\delta}_W(i)>nx\big|\xi^{\delta}_W(0)>nx\right\}\\
&= \frac{\mathbb{P}\left\{\max_{m\leqslant|i|\leqslant r_n}\xi^{\delta}_W(i)\leqslant nx,\xi^{\delta}_W(0)>nx\right\}}{\mathbb{P}\left\{\xi^{\delta}_W(0)>nx\right\}}\\
&= \frac{1-e^{-\mathbb{E}\left\{\max(X^{\delta}(0),\max_{|i|\in\{m,\ldots,r_n\}}X^{\delta}(i))\right\}\frac{1}{nx}}-\left[1-e^{-\mathbb{E}\left\{\max_{|i|\in\{m,\ldots,r_n\}}X^{\delta}(i)\right\}\frac{1}{nx}}\right]}{1-e^{-\frac{1}{nx}}}\\
&\sim nx\Big[1-\big[1-\tfrac{1}{nx}\mathbb{E}\left\{\max\big(1,\max_{|i|\in\{m,\ldots,r_n\}}X^{\delta}(i)\big)\right\}\big]-\Big(1-\big[1-\tfrac{1}{nx}\mathbb{E}\left\{\max_{|i|\in\{m,\ldots,r_n\}}X^{\delta}(i)\right\}\big]\Big)\Big]\\
&\sim \mathbb{E}\left\{\left(1-\max_{|i|\in\{m,\ldots,r_n\}}X^{\delta}(i)\right)_+\right\},
\end{aligned}$$

where we used the fact that as in (4.1), the condition $\lim_{n\to\infty}r_n=\lim_{n\to\infty}\frac{n}{r_n}=\infty$ implies

$$\lim_{n\to\infty}\frac{1}{n}\mathbb{E}\left\{\max\big(X^{\delta}(0),\max_{|i|\in\{m,\ldots,r_n\}}X^{\delta}(i)\big)\right\}=0,$$

and

$$\lim_{n\to\infty}\frac{1}{n}\mathbb{E}\left\{\max_{|i|\in\{m,\dots,r_n\}}X^{\delta}(i)\right\}=0.$$

Consequently,

$$\begin{aligned}\lim_{m\to\infty}\limsup_{n\to\infty}\mathbb{P}\left\{\max_{m\leqslant|i|\leqslant r_n}\xi_W^{\delta}(i)>nx\Big|\xi_W^{\delta}(0)>nx\right\}&=\\ =\lim_{m\to\infty}\limsup_{n\to\infty}\Big[1-\mathbb{E}\left\{\big(1-\max_{|i|\in\{m,\dots,r_n\}}X^{\delta}(i)\big)_+\right\}\Big]&\\ =1-\lim_{m\to\infty}\mathbb{E}\left\{\big(1-\max_{|i|\in\mathbb{Z},i\geqslant m}X^{\delta}(i)\big)_+\right\}&\\ =0,&\end{aligned}$$

where we used the assumption (1.10). Hence Condition 2.1 holds.

In light of [5, Proposition 4.2] we have that Condition 2.1 implies (1.10). Moreover, since

$$\mathbb{P}\left\{\xi_W(0)>n\right\}=1-e^{-1/n}\sim\frac{1}{n},\quad n\to\infty$$

[5, Proposition 4.2] and Lemma 1.1 imply

$$\theta_W^{\delta}=\widetilde{\theta_W^{\delta}}=\widehat{\theta_W^{\delta}}>0.$$

Consequently,

$$\begin{aligned}\widehat{\theta_W^{\delta}}&=\mathbb{P}\left\{\sup_{i\geqslant1}Y^{\delta}(i)\leqslant1\right\}\\ &=\lim_{n\to\infty}\mathbb{P}\left\{\mathcal{P}\sup_{n\geqslant i\geqslant1}X^{\delta}(i)\leqslant1\right\}\\ &=\lim_{n\to\infty}\mathbb{E}\left\{\big(1-\sup_{n\geqslant i\geqslant1}X^{\delta}(i)\big)_+\right\}\\ &=\mathbb{E}\left\{\big(1-\sup_{i\geqslant1}X^{\delta}(i)\big)_+\right\}\\ &=\mathbb{E}\left\{\sup_{i\geqslant0}X^{\delta}(i)-\sup_{i\geqslant1}X^{\delta}(i)\right\}\in(0,1],\end{aligned}\tag{4.2}$$

where the second last expression follows from the monotone convergence theorem. In fact, the above claim readily follows also from [5][Remark 4.7]. Further from (4.2) we obtain

$$\lim_{n\to\infty}\mathbb{P}\left\{\mathcal{P}\sup_{n\geqslant i\geqslant1}X^{\delta}(i)\leqslant1\right\}=\lim_{n\to\infty}\mathbb{P}\left\{\sup_{n\geqslant i\geqslant1}\big(\ln\mathcal{P}+\ln X^{\delta}(i)\big)\leqslant0\right\}$$

$$\begin{aligned} &= \lim_{n\to\infty} \mathbb{P}\left\{ \sup_{n\geqslant i\geqslant 1} \left(\mathcal{E} + W^{\delta}(i)\right) \leqslant 0 \right\} \\ &= \mathbb{P}\left\{ \sup_{i\geqslant 1} \left(\mathcal{E} + W^{\delta}(i)\right) \leqslant 0 \right\}, \end{aligned}$$

with $\mathcal{E} = \ln \mathcal{P}$ a unit exponetial random variable independent of X.

Next, (2.3) follows from [45][Eq. (16)]. Since further we assume (1.2), then (2.3) implies

$$\mathcal{H}_W^{\delta} \in (0, 1/\delta) \tag{4.3}$$

for any $\delta > 0$, establishing thus the proof. □

Proof of Theorem 2.2: By our assumption for all large k

$$\frac{\sigma^2(\delta k)}{8} > \ln(\delta k)^a.$$

Consequently, by (3.8) we have for all δ large and some $a > 1$

$$\mathcal{H}_W^0 \geqslant \mathcal{H}_W^{\delta} \quad \geqslant \quad \frac{1}{\delta}\left(1 - \sum_{k=1}^{\infty} e^{-\frac{\sigma^2(\delta k)}{8}}\right) \geqslant \frac{1}{\delta}\left(1 - \frac{1}{\delta^a}\sum_{k=1}^{\infty}\frac{1}{k^a}\right) > 0.$$

Hence the proof is complete. □

Proof of Proposition 3.1: Since $B_\alpha(0) = 0$ almost surely, in view of (2.6) (see also [19][Proposition 4]) we obtain

$$\lim_{\delta\to 0} \delta^{-1}\mathbb{P}\left\{\forall_{k\in\mathbb{Z}\setminus\{0\}} B_\alpha(\delta k) \leqslant |\delta k|^\alpha/\sqrt{2}\right\} = \mathcal{H}_{B_\alpha}.$$

Moreover, by the self-similarity of B_α, we have

$$\begin{aligned} \mathbb{P}\left\{\forall_{k\in\mathbb{Z}\setminus\{0\}} B_\alpha(\delta k) \leqslant |\delta k|^\alpha/\sqrt{2}\right\} &= \mathbb{P}\left\{\forall_{k\in\mathbb{Z}\setminus\{0\}} |\delta k|^\alpha B_\alpha\left(\frac{1}{\delta k}\right) \leqslant |\delta k|^\alpha/\sqrt{2}\right\} \\ &= \mathbb{P}\left\{\forall_{k\in\mathbb{Z}\setminus\{0\}} B_\alpha\left(\frac{1}{\delta k}\right) \leqslant 1/\sqrt{2}\right\} \\ &= \mathbb{P}\left\{\forall_{k\in\mathbb{Z}\setminus\{0\}} B_\alpha\left(\frac{1}{k}\right) \leqslant \delta^{\alpha/2}/\sqrt{2}\right\}, \end{aligned}$$

hence the proof follows easily. □

Proof of Theorem 3.1: i) The proof is based on a technique developed in Lemma 16 and Corollary 17 in [16] and in Lemma 7 in [48], therefore we omit

some details. For any $\delta > 0$ and T positive integer, using Bonferoni's inequality we have for any process W such that $\mathbb{E}\left\{e^{W(k\delta)}\right\} = 1, k \geqslant 1$

$$
\begin{aligned}
&\mathbb{E}\left\{\sup_{t\in\delta\mathbb{Z}\cap[0,\delta T]} e^{W}(t)\right\}\\
&= \int_{\mathbb{R}} e^s \mathbb{P}\left\{\sup_{t\in\delta\mathbb{Z}\cap[0,\delta T]} W(t) > s\right\} ds\\
&\geqslant \int_{\mathbb{R}} e^s \mathbb{P}\left\{\exists_{1\leqslant k\leqslant T} W(k\delta) > s\right\} ds\\
&\geqslant \sum_{k=1}^{T} \int_{\mathbb{R}} e^s \mathbb{P}\left\{W(k\delta) > s\right\} ds\\
&\quad - \sum_{k=1}^{T-1}\sum_{l=k+1}^{T} \int_{\mathbb{R}} e^s \mathbb{P}\left\{W(k\delta) > s, W(l\delta) > s\right\} ds\\
&\geqslant \sum_{k=1}^{T} \mathbb{E}\left\{e^{W(k\delta)}\right\} \sum_{k=1}^{T-1}\sum_{l=k+1}^{T} \int_{\mathbb{R}} e^s \mathbb{P}\left\{W(k\delta) + W(l\delta) > 2s\right\} ds\\
&= T - \sum_{k=1}^{T-1}\sum_{l=k+1}^{T} \int_{\mathbb{R}} e^s \mathbb{P}\left\{W(k\delta) + W(l\delta) > 2s\right\} ds \qquad (4.4)\\
&= T - \sum_{k=1}^{T-1}\sum_{l=k+1}^{T} \mathbb{E}\left\{e^{\frac{W(k\delta)+W(l\delta)}{2}}\right\}\\
&= T - \sum_{k=1}^{T-1}\sum_{l=k+1}^{T} e^{-\frac{\sigma^2(\delta|k-l|)}{8}}\\
&\geqslant T - T\sum_{k=1}^{T} e^{-\frac{\sigma^2(\delta k)}{8}},
\end{aligned}
$$

where the last equality follows by the stationary of increments of the random process B. Along the lines of the proof in [14]

$$
\begin{aligned}
\mathcal{H}_W^{\delta} &= \lim_{T\to\infty} \frac{1}{T}\mathbb{E}\left\{\sup_{t\in\delta\mathbb{Z}\cap[0,T]} e^{W}(t)\right\}\\
&\geqslant \lim_{T\to\infty} \frac{1}{T}\lfloor T/\delta\rfloor\big(1 - \sum_{k=1}^{\infty} e^{-\frac{\sigma^2(\delta k)}{8}}\big)_+\\
&= \frac{1}{\delta}\big(1 - \sum_{k=1}^{\infty} e^{-\frac{\sigma^2(\delta k)}{8}}\big)_+.
\end{aligned}
$$

ii) In view of (4.4), in order to establish the proof we need to calculate

$$a_{kl} = \int_{\mathbb{R}} e^s \mathbb{P}\left\{W(\delta k) + W(\delta l) > 2s\right\}\, ds.$$

By the independence of the increments, and the fact that $W(\delta l) - W(\delta k) \stackrel{d}{=} W(\delta(l-k))$ we have

$$\begin{aligned} a_{kl} &= \mathbb{E}\left\{e^{\frac{W(\delta k)+W(\delta l)}{2}}\right\} \\ &= \mathbb{E}\left\{e^{W(\delta k)}\right\}\mathbb{E}\left\{e^{\frac{W(\delta l)-W(\delta k)}{2}}\right\} \\ &= \mathbb{E}\left\{e^{W(\delta k)}\right\}\mathbb{E}\left\{e^{\frac{W(\delta (l-k))}{2}}\right\} \\ &= \mathbb{E}\left\{e^{\frac{B(\delta (l-k))-\Phi(1)\delta(l-k)}{2}}\right\} \\ &= \exp\left(-\delta(l-k)\lambda\right), \end{aligned}$$

where $\lambda := \frac{1}{2}\Phi(1) - \Phi(1/2) > 0$ by Jensen's inequality and independence and stationarity of increments of the Lévy process B. Consequently, for $N \in \mathbb{N}$ we obtain

$$\begin{aligned} \int_{\mathbb{R}} e^s \mathbb{P}\left\{\sup_{t\in\delta\mathbb{Z}\cap[0,N]} W(t) > s\right\} ds &\geqslant \frac{N}{\delta}\left(1 - \sum_{k=1}^{\infty} e^{-\delta k\lambda}\right) \\ &= \frac{N}{\delta}\frac{1-2\exp\left(-\delta\lambda\right)}{1-\exp\left(-\delta\lambda\right)}, \end{aligned} \tag{4.5}$$

which leads to

$$\mathcal{H}_W^{\delta} \geqslant \frac{1}{\delta}\frac{1-2\exp\left(-\delta\lambda\right)}{1-\exp\left(-\delta\lambda\right)}$$

and thus the proof is complete. □

Proof of (3.11): By (4.5) and letting $\lambda = \frac{1}{2}\Phi(1) - \Phi(1/2) > 0$ we have

$$\begin{aligned} \mathcal{H}_W^0 &\geqslant \lim_{N\to\infty}\frac{1}{N}\int_{\mathbb{R}} e^s \mathbb{P}\left\{\sup_{t\in\delta\mathbb{Z}\cap[0,N]} W(t) > s\right\} ds \\ &\geqslant \frac{1}{\delta}\left(1 - \sum_{k=1}^{\infty} e^{-\delta k\lambda}\right) \\ &\geqslant \frac{1}{\delta}\left(1 - \int_0^{\infty} e^{-\delta x\lambda} dx\right) \end{aligned}$$

$$\begin{aligned} &= \frac{1}{\delta}\left(1-\frac{1}{\delta\lambda}\right) \\ &\geqslant \frac{\lambda}{4} > 0 \end{aligned}$$

establishing the proof. □

Proof of Proposition 3.2: In view of [11] for any $\delta > 0$ and any integer $k \in \mathbb{N}$ we have

$$\mathcal{H}_W^{\delta} \geqslant \mathbb{E}\left\{\frac{\sup_{t\in\delta\mathbb{Z}} e^{W(t)}}{k\delta \sum\limits_{t\in k\delta\mathbb{Z}} e^{W(t)}}\right\},$$

hence choosing $\delta_n = \eta l^{-n}$ with $\eta > 0$ and $l > 1$ some integer and for $k = l^n$ which is clearly integer for any $n \geqslant 1$ we have

$$\begin{aligned} \mathcal{H}_W^{\delta_n} &\geqslant \mathbb{E}\left\{\frac{\sup_{t\in\delta_n\mathbb{Z}} e^{W(t)}}{k\delta_n \sum\limits_{t\in k\delta_n\mathbb{Z}} e^{W(t)}}\right\} \\ &= \mathbb{E}\left\{\frac{\sup_{t\in\delta_n\mathbb{Z}} e^{W(t)}}{\eta \sum\limits_{t\in\eta\mathbb{Z}} e^{W(t)}}\right\} \\ &\to \mathbb{E}\left\{\frac{\sup_{t\in\mathbb{R}} e^{W(t)}}{\eta \sum\limits_{t\in\eta\mathbb{Z}} e^{W(t)}}\right\}, \quad n\to\infty, \end{aligned}$$

where the last limit follows by the monotone convergence theorem and the fact that W has continuous sample paths. Since by the construction $\mathcal{H}_W^{\delta_n}$ is non-decreasing in n, and we assume that $\lim_{\delta\downarrow 0}\mathcal{H}_W^{\delta} = \overline{H}_W$, then the claim follows. □

Acknowledgements. We are indebted to Bojan Basrak and Philippe Soulier for several comments and suggestions. Thanks to a Swiss National Science Foundation grant no. 200021-166274 for partial support. KD also acknowledges partial support by NCN Grant No 2015/17/B/ST1/01102 (2016-2019).

REFERENCES

[1] J. M. P. Albin, *On extremal theory for stationary processes*, Ann. Probab., 18 (1990), pp. 92–128.

[2] J. M. P. Albin, E. Hashorva, L. Ji, and C. Ling, *Extremes and limit theorems for difference of chi-type processes*, ESAIM: P& S, 20, (2016), pp. 349–366.

[3] B. Basrak, R. A. Davis, and T. Mikosch, *Regular variation of GARCH processes*, Stochastic Process. Appl., 99 (2002), pp. 95–115.

[4] B. Basrak, D. Krizmanić, and J. Segers, *A functional limit theorem for dependent sequences with infinite variance stable limits*, Ann. Probab., 40 (2012), pp. 2008–2033.

[5] B. Basrak and J. Segers, *Regularly varying multivariate time series*, Stochastic Process. Appl., 119 (2009), pp. 1055–1080.

[6] S. Berman, *Sojourns and extremes of stationary processes*, Ann. Probab., 10 (1982), pp. 1–46.

[7] P. J. Bickel and M. Rosenblatt, *On some global measures of the deviations of density function estimates*, Ann. Statist., 1 (1973), pp. 1071–1095.

[8] ———, *Two-dimensional random fields*, in Multivariate analysis, III (Proc. Third Internat. Sympos., Wright State Univ., Dayton, Ohio, 1972), Academic Press, New York, 1973, pp. 3–15.

[9] K. Burnecki and Z. Michna, *Simulation of Pickands constants*, Probab. Math. Statist., 22 (2002), pp. 193–199.

[10] R. A. Davis, T. Mikosch, and Y. Zhao, *Measures of serial extremal dependence and their estimation*, Stochastic Process. Appl., 123 (2013), pp. 2575–2602.

[11] K. Dębicki, S. Engelke, and E. Hashorva, *Brown-Resnick processes and Pickands-type constants*, Extremes, DOI: 10.1007/s10687-017-0289-1, (2017).

[12] K. Dębicki, E. Hashorva, L. Ji, and C. Ling, *Extremes of order statistics of stationary processes*, TEST, 24 (2015), pp. 229–248.

[13] K. Dębicki, *Ruin probability for Gaussian integrated processes*, Stochastic Process. Appl., 98 (2002), pp. 151–174.

[14] K. Dębicki, E. Hashorva, L. Ji, and K. Tabiś, *Extremes of vector-valued Gaussian processes: Exact asymptotics*, Stochastic Process. Appl., 125 (2015), pp. 4039–4065.

[15] K. Dębicki and K. M. Kosiński, *An Erdös-Révész type law of the iterated logarithm for order statistics of a stationary Gaussian process*, J. Theor. Probab., (2016).

[16] K. Dębicki, Z. Michna, and T. Rolski, *Simulation of the asymptotic constant in some fluid models*, Stochastic Models, 19 (2003), pp. 407–423.

[17] M. Delorme, A. Rosso, and K. J. Wiese, *Pickands' constant at first order in an expansion around Brownian motion*, https://arxiv.org/abs/1609.07909, (2016).

[18] A. B. Dieker and T. Mikosch, *Exact simulation of Brown-Resnick random fields at a finite number of locations*, Extremes, 18 (2015), pp. 301–314.

[19] A. B. Dieker and B. Yakir, *On asymptotic constants in the theory of extremes for Gaussian processes*, Bernoulli, 20 (2014), pp. 1600–1619.

[20] C. Dombry and Z. Kabluchko, *Ergodic decompositions of stationary max-stable processes in terms of their spectral functions*, Stochastic Process. Appl., DOI: 10.1016/j.spa.2016.10.00 (2017,in press).

[21] P. Doukhan, A. Jakubowski, and G. Lang, *Phantom distribution functions for some stationary sequences*, Extremes, 18 (2015), pp. 697–725.

[22] P. Embrechts, C. Klüppelberg, and T. Mikosch, *Modelling extremal events for insurance and finance*, Springer-Verlag, Berlin, 1997.

[23] S. Engelke and Z. Kabluchko, *Max-stable processes associated with stationary systems of independent Lévy particles*, Stochastic Process. Appl., 125 (2015), pp. 4272–4299.

[24] M. Falk, J. Hüsler, and R.-D. Reiss, *Laws of small numbers: extremes and rare events*, Birkhäuser/Springer Basel AG, Basel, extended ed., 2011.

[25] A. J. Harper, *Bounds on the suprema of Gaussian processes, and omega results for the sum of a random multiplicative function*, Ann. Appl. Probab., 23 (2013), pp. 584–616.

[26] ———, *Pickands' constant H_α does not equal $1/\Gamma(1/\alpha)$, for small α*, Bernoulli, accepted, (2015).

[27] E. Hashorva, *Representations of max-stable processes via exponential tilting*, arXiv:1605.03208v3, (2016).

[28] J. Hüsler, *Extremes of a Gaussian process and the constant H_α*, Extremes, 2 (1999), pp. 59–70.

[29] A. Janssen and J. Segers, *Markov tail chains*, J. Appl. Probab., 51 (2014), pp. 1133–1153.

[30] Z. Kabluchko, M. Schlather, and L. de Haan, *Stationary max-stable fields associated to negative definite functions*, Ann. Probab., 37 (2009), pp. 2042–2065.

[31] Z. Kabluchko and Y. Wang, *Limiting distribution for the maximal standardized increment of a random walk*, Stochastic Process. Appl., 124 (2014), pp. 2824–2867.

[32] D. Krizmanić, *Functional weak convergence of partial maxima processes*, Extremes, 19 (2016), pp. 7–23.

[33] R. Kulik and P. Soulier, *Heavy tailed time series with extremal independence*, Extremes, 18 (2015), pp. 273–299.

[34] M. Marcus, *Upper bounds for the asymptotic maxima of continuous Gaussian processes*, Ann. Math. Statist., 43 (1972), pp. 522–533.

[35] N. M. Markovich, *Modeling clusters of extreme values*, Extremes, 17 (2014), pp. 97–125.

[36] Z. Michna, *On tail probabilities and first passage times for fractional Brownian motion*, Math. Methods Oper. Res., 49 (1999), pp. 335–354.

[37] T. Mikosch and C. Stărică, *Limit theory for the sample autocorrelations and extremes of a GARCH* $(1, 1)$ *process*, Ann. Statist., 28 (2000), pp. 1427–1451.

[38] T. Mikosch and Y. Zhao, *The integrated periodogram of a dependent extremal event sequence*, Stochastic Process. Appl., 125 (2015), pp. 3126–3169.

[39] I. Molchanov and K. Stucki, *Stationarity of multivariate particle systems*, Stochastic Process. Appl., 123 (2013), pp. 2272–2285.

[40] M. Oesting, Z. Kabluchko, and M. Schlather, *Simulation of Brown-Resnick processes*, Extremes, 15 (2012), pp. 89–107.

[41] J. Pickands, III, *Asymptotic properties of the maximum in a stationary Gaussian process*, Trans. Amer. Math. Soc., 145 (1969), pp. 75–86.

[42] V. I. Piterbarg, *Twenty Lectures About Gaussian Processes*, Atlantic Financial Press, London, New York, 2015.

[43] C. Qualls and H. Watanabe, *Asymptotic properties of Gaussian processes*, Ann. Math. Statist., 43 (1972), pp. 580–596.

[44] G. O. Roberts, J. S. Rosenthal, J. Segers, and B. Sousa, *Extremal indices, geometric ergodicity of Markov chains, and MCMC*, Extremes, 9 (2006), pp. 213–229.

[45] J. Segers, *Approximate distributions of clusters of extremes*, Statist. Probab. Lett., 74 (2005), pp. 330–336.

[46] ———, *Rare events, temporal dependence, and the extremal index*, J. Appl. Probab., 43 (2006), pp. 463–485.

[47] J. Segers, Z. Yuwei, and M. Thomas, *Radial-angular decomposition of regularly varying time series in star-shaped metric spaces*, https://arxiv.org/abs/1604.00241, (2016).

[48] Q. Shao, *Bounds and estimators of a basic constant in extreme value theory of Gaussian processes*, Statistica Sinica, 6 (1996), pp. 245–258.

[49] S. A. Stoev, *On the ergodicity and mixing of max-stable processes*, Stochastic Process. Appl., 118 (2008), pp. 1679–1705.

[50] ———, *Max–stable processes: Representations, ergodic properties and statistical applications*, Dependence in Probability and Statistics, Lecture Notes in Statistics 200, Doukhan, P., Lang, G., Surgailis, D., Teyssiere, G. (Eds.), 200 (2010), pp. 21–42.

[51] Y. Wang, *Extremes of q-Ornstein-Uhlenbeck processes*, https://arxiv.org/abs/1609.00338, (2016).

[52] Y. Wang and S. A. Stoev, *On the structure and representations of max-stable processes*,

Adv. in Appl. Probab., 42 (2010), pp. 855–877.

Mathematical Institute, University of Wrocław
pl. Grunwaldzki 2/4,
50-384 Wrocław, Poland
E-mail: Krzysztof.Debicki@math.uni.wroc.pl

University of Lausanne
Bâtiment Extranef, UNIL-Dorigny,
1015 Lausanne, Switzerland
E-mail: Enkelejd.Hashorva@unil.ch

Received on 24.11.2016;
revised version on xx.xx.xxxx